

TECH-GRO™ B-17 BORIC ACID SPRAY™



B-17 BORIC ACID SPRAY™ is a soluble, fine granular product formulated with improved dispersion and wetting characteristics to provide superior foliar boron performance.

GUARANTEED ANALYSIS

Boron (B)	17.0%
Boron derived from boric acid.	

F1132

GENERAL RECOMMENDATIONS

B-17 BORIC ACID SPRAY at a rate of 5.9 lbs. provides the equivalent of 1 lb. of boron. B-17 contains no sodium and is mildly acidic.

DILUTION: Apply the suggested rates of B-17 BORIC ACID SPRAY in the following amounts of water per acre. Minimum dilution rate - 1 lb. per 5 gallons of water per suggested rate. Allow adequate time for complete dissolution.

- Field & Row Crops
- By air 10-20 gals.
- By ground 10 gals. minimum.
- Tree Crops:
- By air 10 gals. minimum.
- By ground concentrate . . . 40-100 gals.
- By ground dilute 100-500 gals.
- In oil sprays, use 1-5 lbs. per 100 gallons.

COMPATIBILITY: This product has been designed to be compatible with most commonly used insecticides, fungicides and nutritional products. However, a compatibility test is recommended for combinations that have not been used previously. Consult your NUTRI-TECH representative for detailed compatibility information.

FOR SPECIFIC CROP RECOMMENDATIONS - SEE SUGGESTED USES ON BACK

WARNING: This product contains Boron and is to be used only where Boron is known to be beneficial and on the crops recommended on this label. Any other use may result in injury to crops.

ADEQUATE AGITATION SHOULD BE MAINTAINED AT ALL TIMES DURING MIXING AND APPLICATION.

KEEP OUT OF REACH OF CHILDREN

WARNING

Suspected of damaging fertility or the unborn child



Wear appropriate personal protective equipment. Avoid breathing dust. Do not ingest this product. Dispose of unused contents, container, and other contaminated wastes in accordance with local, state, and federal regulations. **If exposed or concerned:** Get medical advice or attention.



NET CONTENTS: 25.0 LBS. (11.3 KG.)



TECH-GRO™ B-17 BORIC ACID SPRAY

SUGGESTED USES (RATES PER ACRE)

FOR ALL CROPS - DO NOT EXCEED THE RATE OF 6.5 LBS. PER ACRE PER YEAR OF B-17 IN FOLIAR APPLICATIONS, 12 LBS. PER ACRE PER YEAR IN SOIL APPLICATIONS. SUGGESTED RATES MAY BE APPLIED IN SPLIT APPLICATIONS.

SOIL APPLICATIONS:

For All Crops: 3-12 lbs. per acre as needed.

FOLIAR APPLICATIONS:

Alfalfa, Hay, & Other Forage Crops - Apply 0.5-3 lbs. over spring growth, repeat after each cutting.

Almonds, Peaches, Plums, Cherries, & Other Stone Fruits - Apply 1-5 lbs. pre-bloom and at 3-4 week intervals as needed post-bloom. (SEE BELOW)

Apples, Pears - Apply 1-6.25 lbs. at 2-4 week intervals starting pre-bloom. (SEE BELOW)

Barley, Oats, Rice, Wheat & Other Small Grains - Apply 0.5-3 lbs. at tillering, repeat at early boot if necessary.

Beans, Peas, & related Legumes - Apply 0.5-3 lbs. at early bud, repeat after bloom if necessary.

Cabbage, Cauliflower, Broccoli, Brussels Sprouts & Other Cole Crops - Apply 0.5-3 lbs. as needed after thinning or transplant.

Carrots, Onions, Radishes, Beets & Other Vegetable Root Crops - Apply 0.5-3 lbs. when top growth is 4-8" high, repeat as necessary.

Citrus, Avocados - Apply 1-5 lbs. in spring, summer, and fall sprays.

Corn, Field & Sweet - Apply 0.5-3 lbs. when 10-20" high, repeat in 2-4 weeks as necessary.

Cotton - Apply 0.5-3 lbs. at first square, repeat in 2-4 weeks as necessary.

Cucumbers, Melons, Squash & Other Cucurbits - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.

Grapes - Apply 0.5-3 lbs. dormant or as needed pre-bloom to post-bloom.

Hops - Apply 0.5-3 lbs. as needed in cover sprays.

Lettuce, Celery, Spinach & Other Green Leafy Vegetables - Apply 0.5-3 lbs. as needed after thinning and transplant.

Mint - Apply 0.5-3 lbs. over new growth, repeat as needed.

Peanuts - Apply 0.5-3 lbs. at early bloom and just prior to pegging.

Pecans, & Other Deciduous Nut Trees - Apply 1-5 lbs. at 3-4 week intervals starting at bud break.

Ornamentals, Trees & Shrubs - Apply 0.5-3 lbs. at 4-6 week intervals as needed.

Potatoes - Apply 0.5-3 lbs. at tuber initiation, repeat in 2-4 weeks as necessary.

Raspberries, Blackberries, Blueberries, Cranberries & Other Small Fruit - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary

Soybeans, Seed Crops - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.

Sugarbeets - Apply 0.5-3 lbs. at 4th-6th leaf, repeat in 2-4 weeks as necessary.

Strawberries - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.

Tomatoes, Peppers, Eggplant & Okra - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.

Walnuts - Apply 1-5 lbs. in blight sprays.

For apples, pears, and all deciduous stone fruits, including almonds, B-17 may be applied in delayed dormant programs and post harvest sprays as per local recommendations.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used according to directions under normal conditions on the crops specified. Because of widely varying use conditions, it is impossible to eliminate all risks even when the label directions are followed. Therefore, the manufacturer makes no other warranties, expressed or implied. The buyer assumes all risks associated with the use of this product. In the event of damage resulting from a breach of warranty, the buyer agrees to accept a refund of the purchase price as full discharge of the manufacturer's liability.

Information regarding the contents and levels of metals in this product is available on the internet at

<http://www.aapfco.org/metals.html>

2.1

www.NutrientTechnologies.com