



GUARANTEED ANALYSIS

Boron (B) 17.0%

Derived From: Boric Acid **F1132**

Uniformly apply the suggested rates in the following amounts of water per acre.

COMPATABILITY: This product is compatible with most crop protection materials. A compatibility test is recommended when using new combinations. Consult a Nutrient TECH™ representative for additional information.

B-17 Boric Acid Spray at a rate of 5.9 lbs. provides the equivalent of 1 lb. of Boron. B-17 contains no sodium and is mildly acidic.

Minimum dilution rate - 1 lb. per 5 gallons of water per suggested rate. Allow adequate time for complete dissolution.

Field & Row Crops:

- By air 10-20 gals.
- By ground 10 gals. minimum

Tree Crops:

- By air 10 gals. minimum
- By ground concentrate 40-100 gals.
- By ground dilute 100-500 gals.
- In oil sprays, use 1-5 lbs per 100 gallons.

Do not apply with phosphate or sulfate fertilizer products. Any previously applied phosphate or sulfate fertilizer should be thoroughly flushed from fertigation equipment prior to application of this product. Apply the product slowly into fertigation equipment in the early part of the water application.

MAINTAIN ADEQUATE AGITATION AT ALL TIMES DURING MIXING AND APPLICATION.

WARNING: This product contains Boron (B) and is to be used only where Boron (B) is known to be beneficial and on the crops recommended on this label. The use of the fertilizing material on any crops other than those recommended may result in serious injury to the crop(s).

WARNING

Suspected of damaging fertility or the unborn child



Wear appropriate personal protective equipment. Avoid breathing dust. Do not ingest this product. Dispose of unused contents, container, and other contaminated wastes in accordance with local, state, and federal regulations. If exposed or concerned: Get medical advice or attention.

KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED

AVOID BREATHING DUST



Warning: Cancer and Reproductive Harm
www.P65Warnings.ca.gov

RECOMMENDED USES (RATES PER ACRE)

FOR ALL CROPS: Do not exceed the rate of 6.5 lbs. per acre per year of B-17 in foliar applications, 12 lbs. per acre per year in soil applications. Suggested rates may be applied in split applications.

FOLIAR APPLICATIONS:

Alfalfa, Hay & Other Forage Crops - Apply 0.5-3 lbs. over spring growth, repeat after each cutting.

Almonds, Peaches, Plums, Cherries & Other Stone Fruits - Apply 1-5 lbs. pre-bloom and at 3-4 week intervals as needed post-bloom. (SEE BELOW)

Apples, Pears - Apply 1-6.25 lbs. at 2-4 week intervals starting pre-bloom. (SEE BELOW)

Barley, Oats, Rice, Wheat & Other Small Grains - Apply 0.5-3 lbs. at tillering, repeat at early boot if necessary.

Beans, Peas, & Related Legumes - Apply 0.5-3 lbs. at early bud, repeat after bloom if necessary.

Cabbage, Cauliflower, Broccoli, Brussels Sprouts & Other Cole Crops - Apply 0.5-3 lbs. as needed after thinning or transplant.

Carrots, Onions, Radishes, Beets & Other Vegetable Root Crops - Apply 0.5-3 lbs. when top growth is 4-8" high, repeat as necessary.

Citrus, Avocados - Apply 1-5 lbs. in spring, summer and fall sprays.

Corn, Field & Sweet - Apply 0.5-3 lbs. when 10-20" high, repeat in 2-4 weeks as necessary.

Cotton - Apply 0.5-3 lbs. at first square, repeat in 2-4 weeks as necessary.

Cucumbers, Melons, Squash & Other Cucurbits - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.

Grapes - Apply 0.5-3 lbs. dormant or as needed pre-bloom to post-bloom.

Hops - Apply 0.5-3 lbs. as needed in cover sprays.

B-17™

Boric Acid Spray

Lettuce, Celery, Spinach & Other Green Leafy Vegetables - Apply 0.5-3 lbs. as needed after thinning and transplant.

Mint - Apply 0.5-3 lbs. over new growth, repeat as needed.

Peanuts - Apply 0.5-3 lbs. at early bloom and just prior to pegging.

Pecans & Other Deciduous Nut Trees - Apply 1-5 lbs. at 3-4 week intervals starting at bud break.

Ornamentals, Trees & Shrubs - Apply 0.5-3 lbs. at 4-6 week intervals as needed.

Potatoes - Apply 0.5-3 lbs. at tuber initiation, repeat in 2-4 weeks as necessary.

Raspberries, Blackberries, Blueberries, Cranberries & Other Small Fruit - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.

Soybeans, Seed Crops - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.

Sugarbeets - Apply 0.5-3 lbs. at 4th-6th leaf, repeat in 2-4 weeks as necessary.

Strawberries - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.

Tomatoes, Peppers, Eggplant & Okra - Apply 0.5-3 lbs. pre-bloom, repeat in 2-4 weeks as necessary.

Walnuts - Apply 1-5 lbs. in blight sprays.

B-17: For apples, pears and all deciduous stone fruits, including almonds, B-17 may be applied in delayed dormant programs and post harvest sprays as per local recommendations.

B-17™

Boric Acid Spray

SOIL APPLICATIONS:

For all crops: Apply 3-12 lbs per acre as needed.

Store in a cool, dry, well-ventilated area away from heat, combustible and incompatible materials. Dispose of unused contents, container, and other contaminated wastes in accordance with local, state, and federal law.

NOTICE: Information regarding the contents and levels of metals in this product is available on the Internet at <https://www.aapfco.org/metals.html>

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY: Nutrient Technologies, Inc. ("NTI") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, NTI makes no other warranties and disclaims all other warranties. See full warranty at www.nutrient.tech/ag-products

B-17 is a trademark of Nutrient Technologies, Inc.
Nutrient TECH and Nutrient.TECH wordmark and stylized logo are trademarks of Nutrient Technologies, Inc.

SPECIMEN LABEL