

GUARANTEED ANALYSIS

Total Nitrogen (N)	9.0%
9.0% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	3.0%
Soluble Potash (K ₂ O)	7.0%

Derived from: Urea, Phosphoric Acid, Potassium Hydroxide.

CONTAINS NON-PLANT FOOD INGREDIENTS:

Humic Acids (derived from Leonardite)..... 2.5%

10.55 lbs/gal at 68°F **F1132**

KEEP OUT OF REACH OF CHILDREN

! WARNING: Causes eye and skin irritation

Wear protective clothing, gloves, eye and face protection. Do not eat, drink or smoke when using this product. Wash hands thoroughly after handling. Remove and wash contaminated clothing before reuse. Avoid breathing vapor, mist or dust from this product. Not for human or animal consumption.

If in eyes: Rinse thoroughly with water for several minutes. Continue rinsing and get medical attention if eye irritation persists.

If on skin: Wash with plenty of soap and water.

If swallowed: Rinse mouth and get medical attention if you feel unwell.

SHAKE WELL BEFORE USING
MAINTAIN ADEQUATE AGITATION AT ALL TIMES DURING MIXING AND APPLICATION.



Warning: Cancer and Reproductive Harm
www.P65Warnings.ca.gov

Humic Acids May Aid in the Uptake of Plant Nutrients

RECOMMENDED USES FOR ROOTERRA

Humic acids may aid in the uptake of plant nutrients.

Uniformly apply the suggested rates in the following amounts of water per acre.

COMPATIBILITY: This product is compatible with most crop protection materials. A compatibility test is recommended when using new combinations. Consult a Nutrient TECH™ representative for additional information.

SOIL APPLICATION RATES (Direct and Fertigation)

VEGETABLES, TOMATO, CUCURBITS and LEAFY

GREENS: 2-5 qts/acre at planting or transplanting through bloom, repeat every 15-21 days

YOUNG PERENNIAL PLANTING (Trees and Vines):

2-5 qts/acre from bud-break every 25-40 days

FRUIT and NUT TREES: 2-5 qts/acre from

budbreak through post harvest

CITRUS, VINES, GRAPES, HOPS: 2-5 qts/acre

from bud-break through post harvest

POTATOES, ROOT VEGETABLES: 2-5 qts/acre

from planting to tuber initiation (row closure)

CORN: 2-3 qt/acre from planting to tasseling, repeat

every 15-21 days

SUGAR BEETS: 2-5 qts/acre from planting to row

closure, repeat every 15-21 days

SEED CROPS, FIELD CROPS: 2-5 qts/acre from

planting to early bloom, repeat every 15-21 days

When applying to soil without a following irrigation

Dilute ROOTERRA to 0.5%-1% concentration with water

ROOT DIPPING/TRANSPLANT SOLUTION

Maximum concentration 0.5% (v/v) with water

FOLIAR APPLICATION (AIR or GROUND)

Maximum concentration 3.0% with water. 3 applications at 1 qt.-4 qts/acre every 10-15 days

VEGETABLES (ALL): at 4th-6th leaf through bloom

YOUNG PERENNIAL PLANTING: on sufficiently developed foliage

POTATOES: at start of tuber initiation (row closure)

CORN, SUGAR BEETS: at 4th-6th leaf

COTTON, BEANS, FIELD CROPS: at 4th-6th leaf

Store in a cool, dry, well-ventilated area away from heat, combustible and incompatible materials. Dispose of unused contents, container, and other contaminated wastes in accordance with local, state, and federal law.

NOTICE: Information regarding the contents and levels of metals in this product is available on the Internet at <https://www.aapfco.org/metals.html>

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY: Nutrient Technologies, Inc. ("NTI") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, NTI makes no other warranties and disclaims all other warranties. See full warranty at www.nutrient.tech/ag-products

ROOTERRA is a trademark of Nutrient Technologies, Inc.

Nutrient TECH and Nutrient.TECH wordmark and stylized logo are trademarks of Nutrient Technologies, Inc.