



KEEP OUT OF REACH OF CHILDREN



WARNING: Toxic to aquatic life with long lasting effects



WARNING: Causes eye and skin irritation

Wear protective clothing, gloves, eye and face protection. Do not eat, drink or smoke when using this product. Wash hands thoroughly after handling. Remove and wash contaminated clothing before reuse. Avoid breathing vapor, mist or dust from this product. Not for human or animal consumption.

If in eyes: Rinse thoroughly with water for several minutes. Continue rinsing and get medical attention if eye irritation persists.

If on skin: Wash with plenty of soap and water.

If swallowed: Rinse mouth and get medical attention if you feel unwell.

SHAKE WELL BEFORE USING
MAINTAIN ADEQUATE AGITATION AT ALL TIMES DURING MIXING AND APPLICATION.



Warning: Cancer and Reproductive Harm
www.P65Warnings.ca.gov

GUARANTEED ANALYSIS

Sulfur (S) 2.5%
 Zinc (Zn)..... 22.0%

Derived from: Zinc Oxysulfate

12.00 lbs/gal at 68°F

F1132

RECOMMENDED USES FOR ZETA ZINC 22

Uniformly apply the suggested rates in the following amounts of water per acre.

COMPATIBILITY: This product is compatible with most crop protection materials. A compatibility test is recommended when using new combinations. Consult a Nutrient TECH™ representative for additional information.

FOR FOLIAR APPLICATION			
	Dilution	Rate	Application Timing
Field, Row & Vegetable Crops	By air: 3-20 gal By ground: 5 gal min	1 pt - 2 qts	Apply every 2-4 weeks as needed, starting pre-bloom
Tree & Vine Crops	By air: 10 gal min By ground dilute: 100-500 gal By ground concentrate: 25-100 gal	1 pt - 4 qts	Apply every 2-4 weeks as needed, starting pre-bloom to post harvest
For Kiwi do not exceed 2 qts per acre per application.			
In oil sprays: use 1-4 pints per 100 gallons		Minimum dilution rate: 1 quart per 3 gallons	

NOTE: Product may leave a visible residue. Do not use within 3-4 weeks of harvest on crops where visible residue is a concern.

FOR TABLE GRAPES: Do not apply within 6-8 weeks of harvest.

NOTE FOR GRAPES: Applications of this product with micronized sulfur products at early cane growth may cause tissue damage to young growth.

NOTE: Some varieties of stone fruits, especially early varieties, are sensitive to Zinc. Foliar application during or followed by damp weather may result in spray injury. Do not make foliar applications on Golden Delicious apples

FOR SOIL APPLICATION			
		Rate	Application Timing
For All Crops		1 qt - 12 qts	Apply to root zone every 1-3 weeks as needed
Do not apply with phosphate or sulfate fertilizer products. Any previously applied phosphate or sulfate fertilizer should be thoroughly flushed from fertigation equipment prior to application of this product. Apply the product slowly into fertigation equipment in the early part of the water application. Do not apply through low flow drip irrigation systems.			

Store in a cool, dry, well-ventilated area away from heat, combustible and incompatible materials. Dispose of unused contents, container, and other contaminated wastes in accordance with local, state, and federal law.

NOTICE: Information regarding the contents and levels of metals in this product is available on the Internet at <https://www.aapfco.org/metals.html>
WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY: Nutrient Technologies, Inc. ("NTI") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, NTI makes no other warranties and disclaims all other warranties. See full warranty at www.nutrient.tech/ag-products

ZETA ZINC 22 is a trademark of Nutrient Technologies, Inc.

Nutrient TECH and Nutrient.TECH wordmark and stylized logo are trademarks of Nutrient Technologies, Inc.