

Concentrated Essential Nutrients for Plant Nutrition

KEEP OUT OF REACH OF CHILDREN

NOTICE: Do not store this product at temperatures bellow 40°F as precipitates may form that are difficult to dissolve. DO NOT expose to temperatures bellow 32°F.

WARNING: This product contains Boron (B) and is to be used only where Boron (B) is known to be beneficial and on the crops recommended on this label. The use of the fertilizing material on any crops other than those recommended may result in serious injury to the crop(s).



WARNING: Causes eye and skin irritation

Wear protective clothing, gloves, eye and face protection. Do not eat, drink or smoke when using this product. Wash hands thoroughly after handling. Remove and wash contaminated clothing before reuse. Avoid breathing vapor, mist or dust from this product. Not for human or animal consumption.

If in eyes: Rinse thoroughly with water for several minutes. Continue rinsing and get medical attention if eye irritation persists.

If on skin: Wash with plenty of soap and water.

If swallowed: Rinse mouth and get medical attention if you feel unwell.

SHAKE WELL BEFORE USING
MAINTAIN ADEQUATE AGITATION AT ALL TIMES
DURING MIXING AND APPLICATION.



Warning: Cancer and Reproductive Harm
www.P65Warnings.ca.gov

GUARANTEED ANALYSIS

Boron (B) 2.5%

Derived from: Boric Acid, Sodium Borate.

8.00 lbs/gal at 68°F

F1132

RECOMMENDED USES FOR LIQUIBOR

Uniformly apply the suggested rates in the following amounts of water per acre.

COMPATIBILITY: This product is compatible with most crop protection materials. A compatibility test is recommended when using new combinations. Consult a Nutrient TECH™ representative for additional information.

FOR FOLIAR APPLICATION			
	Dilution	Rate	Application Timing
Field, Row & Vegetable Crops	By air: 3-20 gal By ground: 5 gal min	1 pt - 4 qts	Apply every 1-4 weeks as needed, starting pre-bloom
For Asparagus do not exceed 2 qts per acre per application.			
Tree & Vine Crops	By air: 10 gal min By ground dilute: 100-500 gal By ground concentrate: 25-100 gal	1 pt - 4 qts	Apply every 1-4 weeks as needed, starting pre-bloom to post harvest
For Kiwi do not exceed 2 qts per acre per application. For Citrus, Walnuts, Pistachios and other deciduous tree nuts maximum rate 6 qts per acre per application.			
In oil sprays: use 1-4 pints per 100 gallons		Minimum dilution rate: 1 quart per 3 gallons	

NOTE: Product may leave a visible residue. Do not use within 3-4 weeks of harvest on crops where visible residue is a concern. FOR TABLE GRAPES: Do not apply within 6-8 weeks of harvest.

NOTE FOR GRAPES: Applications of this product with micronized sulfur products at early cane growth may cause tissue damage to young growth.

FOR SOIL APPLICATION			
	Dilution	Rate	Application Timing
Field, Row, Vegetable, Tree & Vine Crops	1 quart per 3 gallons	1 qt - 12 qts	Apply to root zone every 1-3 weeks as needed
Do not apply with phosphate or sulfate fertilizer products. Any previously applied phosphate or sulfate fertilizer should be thoroughly flushed from fertigation equipment prior to application of this product. Apply the product slowly into fertigation equipment in the early part of the water application.			

Store in a cool, dry, well-ventilated area away from heat, combustible and incompatible materials. Dispose of unused contents, container, and other contaminated wastes in accordance with local, state, and federal law.

NOTICE: Information regarding the contents and levels of metals in this product is available on the Internet at <https://www.aapfco.org/metals.html>
WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY: Nutrient Technologies, Inc. ("NTI") warrants that this Product conforms to the specifications on this label. To the extent consistent with applicable law, NTI makes no other warranties and disclaims all other warranties. See full warranty at www.nutrient.tech/ag-products

LIQUIBOR is a trademark of Nutrient Technologies, Inc.

Nutrient TECH and Nutrient.TECH wordmark and stylized logo are trademarks of Nutrient Technologies, Inc.