

Increase Bulking & Specific Gravity



BRIXX 0-22-24

- Phosphorus together with CO₂ and water, is the primary nutrient for photosynthesis. It is needed for the formation and movement of sugars and starches.
- Potassium functions to regulate stomates for carbon fixation (photosynthesis) and water loss. It is essential for sugar distribution from the leaves into the tubers.
- Boron has a significant role in formation of strong cell walls, root growth and sugar transport in the plant. Boron nutrition improves overall yield and quality.



Rate: 2-3 quarts/acre

Timing: When petioles are trending down and then every 10-14 days up to 1 week before vine kill

Brix is safe to apply all season long



1. **Concentration:** Drives nutrient analysis up; reduces risk to crops
2. **Compatibility:** Quality, reliable products highly compatible with most mixes
3. **Availability:** Proven nutrient uptake via foliar, root, and soil application

Nutrient	N	P ₂ O ₂	K ₂ O	Mg	Ca	S	Zn	Mn	Fe	Cu	B	Mo	Co
% by weight	0	22	24	-	-	-	-	-	-	-	1.5	-	-

For more information contact your Crop Advisor or visit www.Nutrient.TECH



Scan to download the Crop Nutrient Advisor App Today!



Northern CA & Intermountain West
Ryan Sanderson, CCA PCA
 208-565-6431
RSanderson@Nutrient.TECH

Coastal CA, Southern CA, AZ & HI
Craig Wyatt
 831-809-1588
CWyatt@Nutrient.TECH

Pacific Northwest
Michael Ruttan, PCA
 509-378-5357
MRuttan@Nutrient.TECH

CA Central Valley
Devin Lilles
 559-287-7724
DLilles@Nutrient.TECH